


PACSIN2 Antibody (F53991)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F53991-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F53991-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)


| | |
|---------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human, Mouse |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Protein G affinity chromatography |
| UniProt | Q9UNF0 |
| Localization | Cytoplasmic |
| Applications | Western Blot : 1:500-1:1000 IHC (FFPE) : 1:50-100 Immunofluorescence : 1:100 |
| Limitations | This PACSIN2 antibody is available for research use only. |




Western blot testing of 1) human Daudi, 2) human HL-60, 3) mouse brain and 4) mouse heart lysate with PACSIN2 antibody at 1:1000. Predicted molecular weight: 56 kDa.



Immunofluorescence testing of HeLa cells with PACSIN2 antibody (green) and Actin Ab (red) at 1:100.



IHC testing of FFPE human stomach with PACSIN2 antibody at 1:100. HIER: steamed with pH6 citrate buffer.



IHC testing of FFPE human breast cancer with PACSIN2 antibody at 1:50. HIER: steamed with pH6 citrate buffer.

Description

Protein kinase C and casein kinase substrate in neurons protein 2 is a lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus. May also play a role in the formation of caveolae at the cell membrane. Recruits DNM2 to caveolae, and thereby plays a role in caveola-mediated endocytosis.

Application Notes

Titration of the PACSIN2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 342-371 from the human protein was used as the immunogen for the PACSIN2 antibody.

Storage

Aliquot the PACSIN2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.