


# PAK6 Antibody (F50895)

| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F50895-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F50895-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

#### **Bulk quote request**

| Availability       | 1-3 business days                                      |
|--------------------|--|
| Species Reactivity | Human  |
| Format             | Purified   |
| Clonality          | Polyclonal (rabbit origin)                             |
| Isotype            | Rabbit Ig  |
| Purity             | Purified   |
| UniProt            | Q9NQU5   |
| Applications       | Western Blot: 1:1000                                   |
| Limitations        | This PAK6 antibody is available for research use only. |



Western blot analysis of PAK6 antibody in lysate from transiently transfected COS7 cells. Lane 1: negative control, 2: PAK1, 3: PAK2, 4: PAK4, 5: PAK5, and 6: PAK6-expressing cells.

### **Description**

The PAK6 protein shares a high degree of sequence similarity with p21-activated kinase (PAK) family members. The proteins of this family are Rac/Cdc42-associated Ste20-like Ser/Thr protein kinases, characterized by a highly conserved amino-terminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain. PAK kinases are implicated in the regulation of a number of cellular processes, including cytoskeleton rearrangement, apoptosis and the MAP kinase signaling pathway. PAK6 was found to interact with androgen receptor (AR), which is a steroid hormone-dependent transcription factor that is important for male sexual differentiation and development. The p21-activated protein kinase 6 gene was found to be highly expressed in testis and prostate tissues and the encoded protein was shown to cotranslocate into the nucleus with AR in response to androgen.

## **Application Notes**

Titration of the PAK6 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 116-146 from the human protein was used as the immunogen for this PAK6 antibody.

## **Storage**

Aliquot the PAK6 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.