


# Parkin Antibody / PARK2 (F49620)


Catalog No.	Formulation	Size
F49620-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F49620-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	O60260
Applications	Western Blot : 1:250-1:1000 Flow Cytometry : 1:10-1:50 Immunofluorescence : 1:10-1:50 IHC (Paraffin) : 1:10-1:50
Limitations	This Parkin antibody is available for research use only.




Western blot analysis of lysate from (1) SH-SY5Y cell line and (2) human brain tissue lysate using Parkin antibody at 1:1000. Expected molecular weight: 50-60 kDa with multiple smaller isoforms.



Parkin antibody flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control (left histogram). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.



Confocal immunofluorescent analysis of Parkin antibody with NCI-H460 cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 Phalloidin (red). DAPI was used as a nuclear counterstain (blue).



Parkin antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue.

### **Description**

Parkinson is the second most common neurodegenerative disease after Alzheimers. About 1 percent of people over the age of 65 and 3 percent of people over the age of 75 are affected by the disease. The mutation is the most common cause of Parkinson disease identified to date. The function of Park2 is not well-known; however, it may play a role in the ubiquitin-mediated proteolytic pathway. Mutations in this gene are known to cause autosomal recessive juvenile parkinsonism. Alternative splicing of this gene produces three known products of undetermined function.

## **Application Notes**

Titration of the Parkin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 111-140 from the human protein was used as the immunogen for this Parkin antibody.

#### **Storage**

Aliquot the Parkin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.