


PARP9 Antibody (RQ6263)

| Catalog No. | Formulation | Size |
|-------------|---|--------|
| RQ6263 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request

| Availability | 1-3 business days |
|--------------------|---|
| Species Reactivity | Human, Mouse, Rat |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Affinity purified |
| Buffer | Lyophilized from 1X PBS with 2% Trehalose |
| UniProt | Q8IXQ6 |
| Applications | Western Blot : 1-2ug/ml Direct ELISA : 0.1-0.5ug/ml |
| Limitations | This PARP9 antibody is available for research use only. |



Western blot testing of 1) rat skeletal muscle and 2) mouse skeletal muscle lysate with PARP9 antibody. Predicted molecular weight ~96 kDa.

Description

Poly (ADP-ribosyl)ation is a post-translational modification of proteins mediated by one of the 17 members of the poly(ADP-ribose) polymerases (PARP). PARP-9 belongs to the subfamily of macroPARPs, associating one to three macro domains to the PARP domain. Overexpression of PARP-9 stimulates cell migration in vitro, suggesting a role for PARP-9 in the promotion of malignant B cell migration and dissemination in high risk DLBCL. PARP-9 is also likely a transcription coactivator, its overexpression in B lymphocytes, stimulated by IFN?, inducing the transcription of IFN?-controlled genes.

Application Notes

Optimal dilution of the PARP9 antibody should be determined by the researcher.

Immunogen

A human recombinant partial protein (amino acids R117-E824) was used as the immunogen for the PARP9 antibody.

Storage

After reconstitution, the PARP9 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.