


PAX2 Antibody [clone PAX2/8671R] (V4114)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V4114-100UG | 0.2~mg/ml in 1X PBS with $0.1~mg/ml$ BSA (US sourced), $0.05%$ sodium azide | 100 ug |
| V4114-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4114SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Recombinant RABBIT MONOCLONAL

Bulk quote request

| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG, kappa |
| Clone Name | PAX2/8671R |
| Purity | Protein A/G affinity |
| UniProt | Q02962 |
| Localization | Nucleus |
| Applications | Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT |
| Limitations | This PAX2 antibody is available for research use only. |



IHC staining of FFPE human renal cell carcinoma tissue with PAX2 antibody (clone PAX2/8671R) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Recognizes a protein of 42kDa, which is identified as PAX2. It is a member of the paired box family of transcription factors, which is required for development and proliferation of the kidney, brain, and mullerian organs. PAX2 genes contain a highly conserved DNA sequence within the paired box region, which encodes a DNA-binding domain, enabling PAX proteins to bind the promoters of specific genes to transcriptionally regulate their expression. PAX2 is specifically

expressed in the developing central nervous system, eye, ear, and urogenital tract, and is essential for the development of these organs. In normal adult tissues PAX2 was mainly detected in the urogenital system, including kidney, ureteric epithelium, fallopian tube epithelium, ovary and uterus. In tumors, PAX2 has been detected in renal cell carcinomas, Wilms' tumors, nephrogenic adenomas and papillary serous carcinoma of the ovary. PAX2 has been used as a marker for the identification of renal cell carcinoma and ovarian carcinoma by immunohistochemistry.

Application Notes

Optimal dilution of the PAX2 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein (within amino acids 223-354) from the human protein was used as the immunogen for the PAX2 antibody.

Storage

Aliquot the PAX2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.