


## PAX3 Antibody / Paired box 3 [clone PAX3/4700] (V4956)

Catalog No.	Formulation	Size
V4956-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4956-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4956SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a, kappa
<b>Clone Name</b>	PAX3/4700
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P23760
<b>Localization</b>	Nucleus
<b>Applications</b>	ELISA (Order BSA-free Format For Coating) :
<b>Limitations</b>	This PAX3 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free PAX3 antibody (clone PAX3/4700) as confirmation of integrity and purity.

## Description

Pax genes contain paired domains that share strong homology to genes in *Drosophila* which are involved in programming early development. The product of the PAX3 gene is a DNA-binding protein expressed during early neurogenesis. Pax-3

is a protein containing both a paired domain and a paired-type homeodomain. During early neurogenesis, Pax-3 expression is limited to mitotic cells in the ventricular zone of the developing spinal cord and to distinct regions in the hindbrain, midbrain and diencephalon. In 10-12 day embryos, expression of Pax-3 is also seen in neural crest cells of the developing spinal ganglia, the craniofacial mesectoderm and in limb mesenchyme. Mutations in the MITF and Pax-3 genes, encoding transcription factors, are responsible for Waardenburg syndrome II (WSII) and WSI/WSIII, respectively.

## **Application Notes**

Optimal dilution of the PAX3 antibody should be determined by the researcher.

## **Immunogen**

A synthetic peptide from the C-terminus region of the quail protein was used as the immunogen for the PAX3 antibody.

## **Storage**

Aliquot the PAX3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.