
# PCDH10 Antibody / Protocadherin 10 (F53847)

| Catalog No. | Formulation | Size |
|---|---|---|
| F53847-0.2ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.2 ml |
| F53847-0.05ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.05 ml |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Antigen affinity purified |
| **Host** | Rabbit |
| **Clonality** | Polyclonal (rabbit origin) |
| **Isotype** | Rabbit Ig |
| **Purity** | Antigen affinity |
| **UniProt** | Q9P2E7 |
| **Applications** | Western blot: 1:1000-1:2000 |
| **Limitations** | This PCDH10 antibody is available for research use only. |




Western blot testing of human 1) HepG2, 2) PC-3 and 3) U-87 MG cell lysate with PCDH10 antibody at 1:2000. Predicted molecular weight: 113 kDa.

## Description

Protocadherin-10 is a potential calcium-dependent cell-adhesion protein.

## Application Notes

Titration of the PCDH10 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 128-160 from the human protein was used as the immunogen for the PCDH10 antibody.

## Storage

Aliquot the PCDH10 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.


Copyright © NSJ Bioreagents. All rights reserved.