


## Periostin Antibody / POSTN [clone POSTN/3503] (V8697)


| Catalog No.    | Formulation  | Size   |
|----------------|--|--------|
| V8697-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8697-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug  |
| V8697SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                             | 100 ug |

### Bulk quote request

|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days   |
| <b>Species Reactivity</b> | Human   |
| <b>Format</b>             | Purified  |
| <b>Host</b>               | Mouse   |
| <b>Clonality</b>          | Monoclonal (mouse origin)                                   |
| <b>Isotype</b>            | Mouse IgG2b, kappa  |
| <b>Clone Name</b>         | POSTN/3503  |
| <b>Purity</b>             | Protein G affinity chromatography                           |
| <b>UniProt</b>            | Q15063  |
| <b>Localization</b>       | Secreted, cytoplasm   |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT |
| <b>Limitations</b>        | This Periostin antibody is available for research use only. |




IHC staining of FFPE human colon carcinoma with Periostin antibody (clone POSTN/3503). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placenta with Periostin antibody (clone POSTN/3503).  
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

#### Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Periostin antibody (clone POSTN/3503). These results demonstrate the foremost specificity of the POSTN/3503 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free Periostin antibody (clone POSTN/3503) as confirmation of integrity and purity.

## Description

Periostin (PN, POSTN), also designated osteoblast-specific factor 2 (OSF-2), is a disulfide linked protein originally isolated as an osteoblast-specific factor. Periostin is a secreted protein that binds heparin and functions as a ligand for 1). Periostin contains a typical signal sequence, followed by a cysteine-rich domain, a fourfold repeated domain, which shows homology with the insect protein fasciclin, and a C-terminal domain.

## Application Notes

Optimal dilution of the Periostin antibody should be determined by the researcher.

## Immunogen

Recombinant full-length human protein was used as the immunogen for the Periostin antibody.

## Storage

Store the Periostin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

