


## Pf4 Antibody / Platelet factor 4 (R32449)


Catalog No.	Formulation	Size
R32449	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Mouse
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	Q9Z126
<b>Localization</b>	Cytoplasmic, membranous
<b>Applications</b>	Western Blot : 0.5-1ug/ml IHC (FFPE) : 1-2ug/ml ELISA : 0.1-0.5ug/ml (mouse protein tested); request BSA-free format for coating
<b>Limitations</b>	This Pf4 antibody is available for research use only.



IHC staining of FFPE mouse lung tissue with Pf4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of mouse spleen tissue lysate with Pf4 antibody. Expected molecular weight: ~8/16/32 kDa (monomer/dimer/tetramer).

## Description

Platelet factor 4 (PF4) is a small cytokine belonging to the CXC chemokine family that is also known as chemokine (C-X-C motif) ligand 4 (CXCL4). By in situ hybridization, the CXCL4 gene is mapped to chromosome 4q12-q21. Its major physiologic role appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. As a strong chemoattractant for neutrophils and fibroblasts, PF4 probably has a role in inflammation and wound repair.

## Application Notes

Optimal dilution of the Pf4 antibody should be determined by the researcher.

## Immunogen

Amino acids V30-S105 from the mouse protein were used as the immunogen for the Pf4 antibody.

## Storage

Prior to reconstitution, store at 4oC. After reconstitution, the Pf4 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.