


## Phospho-Histone H2AX Antibody (pS139) (F55063)


Catalog No.	Formulation	Size
F55063-0.1ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.1 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	P16104
<b>Applications</b>	Western Blot : 1:1000-1:2000 Immunofluorescence : 1:200-1:400
<b>Limitations</b>	This Phospho-Histone H2AX antibody is available for research use only.



Immunofluorescent staining of mouse NIH 3T3 cells untreated (left) and treated (right) with Hydroxyurea, using Phospho-Histone H2AX antibody.



Western blot testing of cell lysate from mouse NIH 3T3 cells untreated (left) or treated (right) with Hydroxyurea, using Phospho-Histone H2AX antibody.

## Description

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation. Included within the [Histone H2A antibodies](#) collection, this antibody enables analysis of histone modification patterns and chromatin regulatory mechanisms involving H2A and its variants.

## Application Notes

Titration of the Phospho-Histone H2AX antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

The amino acid sequence surrounding phosphorylated serine 139 was used as the immunogen for the Phospho-Histone H2AX antibody.

## Storage

Aliquot the Phospho-Histone H2AX antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.