


## Phospho-IRAK1 Antibody (S376) (F54105)

Catalog No.	Formulation	Size
F54105-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F54105-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	P51617
<b>Applications</b>	Western Blot : 1:1000
<b>Limitations</b>	This phospho-IRAK1 antibody is available for research use only.



Western blot analysis of lysates from the human HeLa cell line, untreated or treated with IL-1 beta (20ng/ml) + Calyculin A (100nM), using phospho-IRAK1 antibody (upper) or Tubulin Ab (lower).

## Description

Serine/threonine-protein kinase that plays a critical role in initiating innate immune response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways. Is rapidly recruited by MYD88 to the receptor- signaling complex upon TLR activation. Association with MYD88 leads to IRAK1 phosphorylation by IRAK4 and subsequent autophosphorylation and kinase activation. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin-binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6 complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent

degradation. Phosphorylates the interferon regulatory factor 7 (IRF7) to induce its activation and translocation to the nucleus, resulting in transcriptional activation of type I IFN genes, which drive the cell in an antiviral state. When sumoylated, translocates to the nucleus and phosphorylates STAT3. [UniProt]

## Application Notes

The stated application concentrations are suggested starting points. Titration of the phospho-IRAK1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 348-381 from human IRAK1 was used as the immunogen for the phospho-IRAK1 antibody.

## Storage

Aliquot the phospho-IRAK1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.