


Phospho-ULK1 (pSer556) Antibody / Unc-51 like autophagy activating kinase 1 [clone 31U30] (FY13231)

| Catalog No. | Formulation | Size |
|-------------|--|--------|
| FY13231 | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA | 100 ul |


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

| | |
|--------------------|---|
| Availability | 2-3 weeks |
| Species Reactivity | Human, Mouse |
| Format | Liquid |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG |
| Clone Name | 31U30 |
| Purity | Affinity chromatography |
| Buffer | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| UniProt | O75385 |
| Applications | Immunofluorescence : 1:50-1:200 Immunocytochemistry/Western Blot : 1:500-1:2000 |
| Limitations | This Phospho-ULK1 (pSer556) antibody is available for research use only. |



Immunofluorescent analysis using the Phospho-ULK1 (pSer556) antibody (green) at 1:50 dilution.



Western blot analysis of ULK1 using anti-Phospho-ULK1 (pSer556) antibody. Lane 1: human 293T whole cell lysates, Lane 2: human MCF-7 whole cell lysates, Lane 3: human whole cell lysates, Lane 4: rat PC-12 whole cell lysates, Lane 5: mouse NIH/3T3 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Phospho-ULK1 (Ser556) antibody at 1:500 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal was developed using enhanced chemiluminescent. Western blot detection of phospho-ULK1 (Ser556) shows a band at ~150 kDa across multiple cell lines. Although the predicted molecular weight of ULK1 is ~113 kDa, the phosphorylated form commonly migrates higher due to multiple phosphorylation events and other post-translational modifications.

Description

Phospho-ULK1 (pSer556) antibody detects Unc-51 like autophagy activating kinase 1 phosphorylated at serine 556, encoded by the ULK1 gene. ULK1 is a serine/threonine kinase that initiates autophagy in response to nutrient status and energy stress. Phosphorylation at Ser556 is an important regulatory event that integrates upstream signaling pathways to control autophagosome formation. Phospho-ULK1 (pSer556) antibody provides researchers with a highly specific reagent to study the regulation of autophagy and its role in physiology and disease.

ULK1 functions as the mammalian homolog of yeast Atg1, acting at the top of the autophagy hierarchy. Research using Phospho-ULK1 (Ser556) antibody has demonstrated that phosphorylation at serine 556 by AMP activated protein kinase promotes activation of ULK1 under conditions of energy depletion. This modification stimulates autophagy initiation by enabling the ULK1 complex to recruit downstream autophagy machinery. By contrast, mTORC1 signaling suppresses ULK1 through phosphorylation at other sites, highlighting the dynamic control of ULK1 activity by nutrient and energy signals.

Phospho-ULK1 (Ser556) antibody has been used to show that phosphorylation at this site acts as a molecular switch for autophagy induction. In starvation or low energy states, phosphorylation at Ser556 ensures cells adapt by recycling cytoplasmic contents through lysosomal degradation. This process provides metabolic substrates for survival and maintains homeostasis. Inhibiting phosphorylation at Ser556 impairs autophagy, underscoring its essential role in this pathway.

Dysregulation of ULK1 phosphorylation has been linked to human disease. Studies with Phospho-ULK1 (pSer556) antibody have revealed that impaired autophagy contributes to neurodegenerative diseases, metabolic disorders, and cancer. In cancer, altered ULK1 phosphorylation influences tumor cell survival under stress conditions, promoting growth in nutrient-poor environments. In neurodegeneration, reduced ULK1 activation leads to accumulation of toxic proteins. These findings emphasize the clinical importance of monitoring ULK1 phosphorylation states.

Phospho-ULK1 (pSer556) antibody is widely used in western blotting, immunohistochemistry, and immunofluorescence. Western blotting reveals phosphorylation-dependent shifts in ULK1 activity, immunohistochemistry highlights tissue-specific patterns of autophagy activation, and immunofluorescence demonstrates dynamic localization of phosphorylated ULK1 at autophagosome initiation sites. These applications make Phospho-ULK1 (Ser556) antibody an indispensable reagent in autophagy research.

By supplying validated Phospho-ULK1 (pSer556) antibody reagents, NSJ Bioreagents supports research into energy sensing, autophagy regulation, and disease mechanisms. Detection of phosphorylation at Ser556 provides a precise marker of ULK1 activation and autophagy initiation.

Application Notes

Optimal dilution of the Phospho-ULK1 (pSer556) antibody should be determined by the researcher.

Immunogen

A synthesized peptide derived from human Phospho-ULK1 (pS556) was used as the immunogen for the Phospho-ULK1 (pSer556) antibody.

Storage

Store the Phospho-ULK1 (pSer556) antibody at -20oC.