


## PIGC Antibody (F53916)

Catalog No.	Formulation	Size
F53916-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q92535
<b>Applications</b>	Western Blot : 1:1000-2000
<b>Limitations</b>	This PIGC antibody is available for research use only.



Western blot testing of human 1) HL-60, 2) K562, 3) MCF-7, 4) MOLT-4 and 5) SH-SY5Y cell lysate with PIGC antibody at 1:2000. Predicted molecular weight: 34 kDa.

## Description

Phosphatidylinositol N-acetylglucosaminyltransferase subunit C is part of the complex catalyzing the transfer of N-acetylglucosamine from UDP-N-acetylglucosamine to phosphatidylinositol, the first step of GPI biosynthesis.

## Application Notes

Titration of the PIGC antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 81-112 from the human protein was used as the immunogen for the PIGC antibody.

## Storage

Aliquot the PIGC antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.