


## Plakoglobin Antibody / Gamma Catenin [clone CTNG/1483] (V3235)


Catalog No.	Formulation	Size
V3235-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3235-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3235SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	CTNG/1483
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P14923
<b>Localization</b>	Cell surface, cytoplasmic, cell junctions
<b>Applications</b>	Immunohistochemistry (FFPE) : 0.1-0.2ug/ml for 30 min at RT
<b>Limitations</b>	This Plakoglobin antibody is available for research use only.



IHC testing of FFPE human cervical carcinoma with Plakoglobin antibody (clone CTNG/1483). Required HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min.



IHC testing of FFPE human skin with Plakoglobin antibody (clone CTNG/1483).  
Required HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min.

## Description

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Plakoglobin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

Recombinant human protein was used as the immunogen for the Plakoglobin antibody.

## Storage

Store the Plakoglobin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).