


## PRMT4 Antibody / CARM1 (R31387)


Catalog No.	Formulation	Size
R31387	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

### Bulk quote request


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	Q86X55
<b>Localization</b>	Nuclear, cytoplasmic
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells
<b>Limitations</b>	This PRMT4 antibody is available for research use only.



Immunofluorescent staining of FFPE human HeLa cells with PRMT4 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human 293T, 2) human HeLa, 3) human MCF7, 4) human HepG2, 5) rat testis and 6) rat C6 cell lysate with PRMT4 antibody. Predicted molecular weight ~66 kDa.



Flow cytometry testing of fixed and permeabilized human 293T cells with PRMT4 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= PRMT4 antibody.

## Description

CARM1 (coactivator-associated arginine methyltransferase 1), also known as PRMT4, is an enzyme encoded by the CARM1 gene found in human beings, as well as many other mammals. This gene is mapped to 19p13.2. CARM1/PRMT4 is a regulator of cyclin E1 and DHFR mRNA expression. Its main function includes catalyzing the transfer of a methyl group from S-adenosyl-L-methionine to the side chain nitrogens of arginine residues within proteins to form methylated arginine derivatives and S-adenosyl-L-homocysteine. PRMT4 is a secondary coactivator through its association with p160 family (SRC-1, GRIP1, AIB) of coactivators. It is responsible for moving cells toward the inner cell mass in developing blastocysts. This gene also plays an important role in androgen receptors and may play a role in prostate cancer progression.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the PRMT4 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the middle region of human CARM1/PRMT4 (HLAPFTDEQLYMEQFTKA) was used as the immunogen for this PRMT4 antibody (100% homologous in human, mouse and rat).

## Storage

After reconstitution, the PRMT4 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.