


Prolactin Antibody / PRL [clone PRL/4908R] (V9172)

Catalog No.	Formulation	Size
V9172-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9172-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9172SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	PRL/4908R
Purity	Protein A/G affinity
UniProt	IDP01236
Localization	Secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Prolactin antibody is available for research use only.




IHC staining of FFPE human pituitary tissue with Prolactin antibody (clone PRL/4908R).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human pituitary carcinoma tissue with Prolactin antibody (clone PRL/4908R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Negative control: IHC staining of FFPE human brain tissue using Prolactin antibody (clone PRL/4908R) at 2ug/ml in PBS for 30min RT. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Prolactin antibody (clone PRL/4908R) as confirmation of integrity and purity.

Description

Prolactin is a growth factor that is secreted by the anterior pituitary. It is necessary for the proliferation and differentiation of the mammary glands. Prolactin is useful in the classification of pituitary tumors and study of pituitary disease. It also plays a role in the development of mammary cancer, functioning dually as a mitogen and a differentiating agent.

Application Notes

Optimal dilution of the Prolactin antibody should be determined by the researcher.

Immunogen

A portion of amino acids 63-201 was used as the immunogen for the Prolactin antibody.

Storage

Aliquot the Prolactin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.