


Prostatic acid phosphatase Antibody / PAP / PSAP / ACP3 [clone ACPP/4495R] (V5002)

Catalog No.	Formulation	Size
V5002-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5002-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5002SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	ACPP/4495R
Purity	Protein A/G affinity
UniProt	P15309
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This Prostatic acid phosphatase antibody is available for research use only.



IHC staining of FFPE human prostate carcinoma tissue with Prostatic acid phosphatase antibody (clone ACPP/4495R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate carcinoma tissue with Prostatic acid phosphatase antibody (clone ACPP/4495R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Prostatic Acid Phosphatase (PSAP or PAP) is a member of the histidine acid phosphatase family. It is a non-specific phosphatase that is capable of dephosphorylating tyrosine residues as well as phospholipids under mildly acidic conditions. Prostatic specific acid phosphatase (PSAP) is involved in prostate epithelial differentiation. It is highly expressed in prostate, primary and metastatic carcinoma of the prostate. This marker may be helpful in pinpointing the site of origin in cases of metastatic carcinoma of the prostate, and is considered a more sensitive marker than PSA.

Application Notes

Optimal dilution of the Prostatic acid phosphatase antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 1-100) from the human protein was used as the immunogen for the Prostatic acid phosphatase antibody.

Storage

Aliquot the Prostatic acid phosphatase antibody and store frozen at -200C or colder. Avoid repeated freeze-thaw cycles.