


# Proteasome subunit beta type-6 Antibody / PSMB6 (RQ6512)


Catalog No.	Formulation	Size
RQ6512	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat, Monkey
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P28072
Applications	Western Blot : 1-2ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This Proteasome subunit beta type-6 antibody is available for research use only.



Flow cytometry testing of human A431 cells with Proteasome subunit beta type-6 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Proteasome subunit beta type-6 antibody.



Western blot testing of 1) human Caco-2, 2) human HeLa, 3) monkey COS-7, 4) human A549, 5) human Jurkat, 6) human MCF7, 7) rat liver, 8) rat brain, 9) rat lung, 10) rat testis, 11) mouse liver, 12) mouse brain, 13) mouse lung and 14) mouse testis lysate with Proteasome subunit beta type-6 antibody. Predicted molecular weight ~23 kDa.

## **Description**

Proteasome subunit beta type-6 also known as 20S proteasome subunit beta-1 (based on systematic nomenclature) is a protein that in humans is encoded by the PSMB6 gene. The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. The encoded protein is a member of the proteasome B-type family, also known as the T1B family, and is a 20S core beta subunit in the proteasome. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

# **Application Notes**

Optimal dilution of the Proteasome subunit beta type-6 antibody should be determined by the researcher.

## **Immunogen**

An E. coli-derived human protein (amino acids A50-D201) was used as the immunogen for the Proteasome subunit beta type-6 antibody.

## **Storage**

After reconstitution, the Proteasome subunit beta type-6 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.