


## PRPF38A Antibody / Pre-mRNA-splicing factor 38A (F55049)


Catalog No.	Formulation	Size
F55049-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F55049-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	Q8NAV1
<b>Localization</b>	Nuclear
<b>Applications</b>	Flow Cytometry : 1:10-1:50 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:50-1:100 Western Blot : 1:500-1:1000
<b>Limitations</b>	This PRPF38A antibody is available for research use only.




Western blot testing of human 1) HeLa and 2) Jurkat cell lysate with PRPF38A antibody.  
Predicted molecular weight ~37 kDa.



Western blot testing of human 1) MDA-MB-231, 2) CCRF-CEM and 3) HL60 cell lysate with PRPF38A antibody. Predicted molecular weight ~37 kDa.



IHC testing of FFPE human cervical carcinoma tissue with PRPF38A antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Flow cytometry testing of human HepG2 cells with PRPF38A antibody; Blue=isotype control, Green= PRPF38A antibody.

## Description

Pre-mRNA-splicing factor 38A (PR38A) is involved in pre-mRNA splicing as a component of the spliceosome.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the PRPF38A antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 283-312 from the human protein was used as the immunogen for the PRPF38A antibody .

## Storage

Aliquot the PRPF38A antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.