


pS2 Antibody / TFF1 [clone SPM573] (V9082)

Catalog No.	Formulation	Size
V9082-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V9082-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V9082SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V9082IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM573
Purity	Protein G affinity chromatography
UniProt	P04155
Localization	Cytoplasmic, secreted
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This pS2 antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human breast carcinoma stained with pS2 antibody (SPM573).

Description

It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. It is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.

Application Notes

The optimal dilution of the pS2 antibody for each application should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 57-84 (CFDDTVRGVPWCFYPNTIDVPPEEECEF) from the human protein were used as the immunogen for this pS2 antibody.

Storage

Store the pS2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).