
PTBP1 Antibody (RQ5530)


Catalog No.	Formulation	Size
RQ5530	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P26599
Applications	Western blot: 0.25-0.5ug/ml Immunohistochemistry (FFPE): 1-2ug/ml Immunofluorescence: 2-4ug/ml Flow cytometry: 1-3ug/million cells Direct ELISA: 0.1-0.5ug/ml
Limitations	This PTBP1 antibody is available for research use only.




Western blot testing of human 1) placenta, 2) K562, 3) A431, 4) A549, 5) Caco-2, 6) HEK293 and 7) U-2 OS lysate with PTBP1 antibody. Predicted molecular weight ~57 kDa.




Western blot testing of 1) rat testis, 2) mouse thymus, 3) mouse lung, 4) mouse kidney, 5) mouse testis and 6) mouse SP20 lysate with PTBP1 antibody. Predicted molecular weight ~57 kDa.




Immunofluorescent staining of FFPE U-2 OS cells with PTBP1 antibody (green) and DAPI (blue). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.




Flow cytometry testing of human 293T cells with PTBP1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= PTBP1 antibody.




IHC staining of FFPE human colon with PTBP1 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer with PTBP1 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.




IHC staining of FFPE mouse kidney with PTBP1 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE mouse intestine with PTBP1 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE rat kidney with PTBP1 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE rat intestine with PTBP1 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.

Description

Polypyrimidine tract-binding protein 1 is a protein that in humans is encoded by the PTBP1 gene. It is mapped to 19p13.3. This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA-binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has four repeats of quasi-RNA recognition motif (RRM) domains that bind RNAs. This protein binds to the intronic polypyrimidine tracts that requires pre-mRNA splicing and acts via the protein degradation ubiquitin-proteasome pathway. It may also promote the binding of U2 snRNP to pre-mRNAs. This protein is localized in the nucleoplasm and it is also detected in the perinucleolar structure. Alternatively spliced transcript variants encoding different isoforms have been described.

Application Notes

Optimal dilution of the PTBP1 antibody should be determined by the researcher.

Immunogen

A human recombinant protein (amino acids M1-A504) was used as the immunogen for the PTBP1 antibody.

Storage

After reconstitution, the PTBP1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.