
PTBP2 Antibody [clone 13D11] (RQ6292)


Catalog No.	Formulation	Size
RQ6292	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a
Clone Name	13D11
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9UKA9
Localization	Nuclear
Applications	Western blot: 1-2ug/ml Immunohistochemistry (FFPE): 2-5ug/ml Immunofluorescence: 5ug/ml Flow cytometry: 1-3ug/million cells
Limitations	This PTBP2 antibody is available for research use only.




IHC staining of FFPE mouse testis with PTBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat testis with PTBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human gallbladder adenocarcinoma tissue with PTBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human U-2 OS cells with PTBP2 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human SH-SY5Y, 2) rat testis and 3) mouse testis lysate with PTBP2 antibody. Predicted molecular weight ~57 kDa.



Flow cytometry testing of human ThP-1 cells with PTBP2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= PTBP2 antibody.

Description

Polypyrimidine tract binding protein 2, also known as PTBP2, is a protein which in humans is encoded by the PTBP2 gene. It is mapped to 1p21.3. The protein encoded by this gene binds to intronic polypyrimidine clusters in pre-mRNA molecules and is implicated in controlling the assembly of other splicing-regulatory proteins. This protein is very similar to the polypyrimidine tract binding protein (PTB) but most of its isoforms are expressed primarily in the brain. Alternative splicing results in multiple transcript variants.

Application Notes

Optimal dilution of the PTBP2 antibody should be determined by the researcher.

Immunogen

A human recombinant partial protein (amino acids M1-A504) was used as the immunogen for the PTBP2 antibody.

Storage

After reconstitution, the PTBP2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.