


PTPRR Antibody / Protein tyrosine phosphatase receptor type R [clone 29P99] (FY12004)

Catalog No.	Formulation	Size
FY12004	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium	100 ul
	azide and 50% glycerol, 0.4-0.5mg/ml BSA.	


Recombinant RABBIT MONOCLONAL

Bulk quote request


Availability	2-3 weeks	
Species Reactivity	Human, Mouse, Rat	
Format	Liquid	
Clonality	Recombinant Rabbit Monoclonal	
Isotype	Rabbit IgG	
Clone Name	29P99	
Purity	Affinity-chromatography	
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.	
UniProt	Q15256	
Localization	Perinuclear, cell membrane	
Applications	Western Blot : 1:500-1:2000 Immunohistochemistry : 1:50-1:200	
Limitations	This PTPRR antibody is available for research use only.	



Western blot analysis of PTPRR using anti-PTPRR antibody. Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PTPRR antibody at 1:500 overnight at 4oC, then washed with TBS-10%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal was developed using an ECL Plus Western Blotting Substrate. A specific band was detected for PTPRR at approximately 70 kDa. The expected band size for PTPRR is at 70 kDa.



IHC analysis of PTPRR using anti-PTPRR antibody. PTPRR was detected in a paraffinembedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-PTPRR antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Western blot analysis of PTPRR using anti-PTPRR antibody. Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PTPRR antibody at 1:500 overnight at 4oC, then washed with TBS-10%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal was developed using an ECL Plus Western Blotting Substrate. A specific band was detected for PTPRR at approximately 70 kDa. The expected band size for PTPRR is at 70 kDa.

Description

PTPRR antibody recognizes protein tyrosine phosphatase receptor type R, an enzyme that belongs to the receptor-type protein tyrosine phosphatase family. This phosphatase regulates phosphorylation states within signaling cascades, modulating pathways critical for cell differentiation, adhesion, and neuronal communication. PTPRR is expressed predominantly in the brain, where it interacts with and dephosphorylates mitogen-activated protein kinases (MAPKs). By controlling MAPK activity, PTPRR influences neuronal development and synaptic plasticity, making it an important regulator in the central nervous system.

Studies using PTPRR antibody have associated this protein with neurological disorders. Alterations in PTPRR signaling may contribute to diseases such as Parkinson's disease, Alzheimer's disease, and schizophrenia due to its involvement in regulating MAPK signaling balance. PTPRR has also been implicated in tumorigenesis, as phosphatases can act as either tumor suppressors or facilitators of oncogenic signaling depending on cellular context. Research continues to explore whether PTPRR expression patterns could serve as biomarkers or therapeutic targets.

Antibodies against PTPRR have been validated for applications including western blot, immunohistochemistry, and immunofluorescence. These reagents allow investigators to detect protein distribution in neuronal tissue or evaluate changes in phosphorylation-dependent signaling under experimental conditions. Clone-based versions provide high specificity, ensuring consistent results in both in vitro and in vivo studies.

NSJ Bioreagents provides this PTPRR antibody for research use, supporting investigations into brain development, signaling regulation, and disease mechanisms. Alternate names include protein tyrosine phosphatase receptor type R antibody, PTP-SL antibody, PTPRQ antibody, and PTPR-SL antibody.

Application Notes

Optimal dilution of the PTPRR antibody should be determined by the researcher.

Immunogen

A synthesized peptide derived from human PTPRR was used as the immunogen for the PTPRR antibody.

Storage

