



# PU.1 Antibody / SPI1 [clone TFPU1-1] (V3882)

Catalog No.	Formulation	Size
V3882-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3882-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V38825SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	TFPU1-1
Purity	Protein G affinity chromatography
UniProt	P17947
Localization	Nuclear
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT Immunofluorescence: 1-2ug/ml Western Blot: 1-2ug/ml
Limitations	This PU.1 antibody is available for research use only.






IHC staining of FFPE human lymph node with PU.1 antibody (clone TFPU1-1). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC staining of FFPE human Hodgkin's lymphoma with PU.1 antibody (clone TFPU1-1). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



Immunofluorescent staining of PFA-fixed human Ramos cells with PU.1 antibody (green, clone TFPU1-1) and Phalloidin (red).

### **Description**

Binds to the PU-box, a purine-rich DNA sequence (5'-GAGGAA-3') that can act as a lymphoid-specific enhancer. This protein is a transcriptional activator that may be specifically involved in the differentiation or activation of macrophages or B-cells. Also binds RNA and may modulate pre-mRNA splicing (By similarity). [UniProt]

#### **Application Notes**

The stated application concentrations are suggested starting points. Titration of the PU.1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 16-170 from the human protein was used as the immunogen for the PU.1 antibody.

#### **Storage**

Store the PU.1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).