

# RALGAPB Antibody / Ral GTPase-activating protein subunit beta (FY12459)

Catalog No.	Formulation	Size
FY12459	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml	100 ug

### **Bulk quote request**

Availability	1-2 days
Species Reactivity	Human, Mouse, Rat
Format	Lyophilized
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Immunogen affinity purified
Buffer	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
UniProt	Q86X10
Applications	Western Blot : 0.25-0.5ug/ml Immunohistochemistry : 2-5ug/ml ELISA : 0.1-0.5ug/ml
Limitations	This RALGAPB antibody is available for research use only.

## **Description**

RALGAPB antibody detects Ral GTPase-activating protein subunit beta, a large scaffolding component of the RalGAP complex that inactivates small GTPases RALA and RALB. The RalGAP complex consists of catalytic (RALGAPA1 or RALGAPA2) and regulatory (RALGAPB) subunits and serves as a negative regulator of Ral signaling, which is involved in membrane trafficking, cell migration, and tumorigenesis. The RALGAPB antibody is an essential reagent for exploring Ralmediated signaling pathways and understanding their impact on oncogenic transformation and vesicular transport.

RALGAPB is encoded by the RALGAPB gene located on human chromosome 20q13.33. The protein is cytoplasmic and interacts with Ral GTPases to stimulate their intrinsic GTP hydrolysis activity, thereby switching them from active (GTP-bound) to inactive (GDP-bound) states. Through this action, RALGAPB contributes to regulation of endocytosis, exocytosis, and actin cytoskeleton dynamics. It also links Ral signaling to insulin-stimulated GLUT4 vesicle trafficking in metabolic tissues.

The RALGAPB antibody is widely used to detect the 185–200 kDa protein in western blot and to visualize cytoplasmic localization in immunofluorescence. RALGAPB forms a heterodimer with RALGAPA1 or RALGAPA2, and the resulting complex controls Ral-dependent regulation of mTOR signaling, autophagy, and vesicle movement. Studies have

implicated RALGAPB in insulin sensitivity, as well as in cancer progression where Ral signaling is hyperactivated. Loss or mutation of RALGAPB leads to sustained Ral activity, promoting cellular proliferation and migration.

Functional studies using this antibody have connected RALGAPB to the modulation of cell junction dynamics and receptor trafficking. It is also implicated in neuronal vesicle regulation and autophagosome maturation. NSJ Bioreagents offers a validated RALGAPB antibody optimized for western blot, immunoprecipitation, and immunofluorescence, enabling researchers to examine its role in signaling, vesicular transport, and metabolic regulation.

#### **Application Notes**

Optimal dilution of the RALGAPB antibody should be determined by the researcher.

#### **Immunogen**

E.coli-derived human RALGAPB recombinant protein (Position: M1-H1025) was used as the immunogen for the RALGAPB antibody.

#### **Storage**

After reconstitution, the RALGAPB antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.