


RBPMs Antibody / RNA-Binding Protein With Multiple Splicing Antibody (RQ6004)


Catalog No.	Formulation	Size
RQ6004	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q93062
Localization	Nuclear, cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry : 2-5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This RBPMs Antibody / RNA-Binding Protein With Multiple Splicing Antibody is available for research use only.




RBPMs Antibody Human Caco-2 Colorectal Cancer Cell FACS. Flow cytometry analysis of fixed and permeabilized human Caco-2 colorectal cancer cells was performed with RBPMs Antibody at 1 ug/million cells. RBPMs staining (blue) shows a clear fluorescence shift compared with the isotype control (green) and blank control (red). This RBPMs Antibody demonstrates intracellular detection of RNA-Binding Protein With Multiple Splicing in human Caco-2 colorectal cancer cells.




RBPMs Antibody Human, Mouse and Rat WB. Western blot analysis was performed with RBPMs Antibody on human Caco-2, A549, RT4 and HepG2 cell lysates (lanes 1-4), rat heart tissue lysate (lane 5) and mouse heart tissue lysate (lane 6). Samples were separated by 12% SDS-PAGE under reducing conditions, and the antibody was used at 0.5 ug/ml overnight at 4oC. Bands are detected at approximately 24 and 28 kDa. Although the calculated molecular weight of RBPMs is approximately 22 kDa, RBPMs has been reported to migrate at approximately 28-30 kDa by Western blot, with multiple bands potentially reflecting splice variants and/or post-translationally modified forms. This RBPMs Antibody demonstrates detection of RNA-Binding Protein With Multiple Splicing across human, mouse and rat samples.




RBPMs Antibody Human Cardiac Muscle IHC. IHC staining of FFPE human cardiac muscle tissue was performed with RBPMs Antibody at 2 ug/ml following heat-induced epitope retrieval in pH 8 EDTA buffer. HRP-conjugated secondary antibody and DAB were used for detection. RBPMs staining is observed in cardiac muscle cells, with predominantly cytoplasmic signal and stronger staining in scattered cells. This RBPMs Antibody demonstrates detection of RNA-Binding Protein With Multiple Splicing in human cardiac muscle tissue.



RBPMs Antibody Human Cardiac Muscle IHC. IHC staining of FFPE human cardiac muscle tissue was performed with RBPMs Antibody at 2 ug/ml following heat-induced epitope retrieval in pH 8 EDTA buffer. HRP-conjugated secondary antibody and DAB were used for detection. RBPMs staining is observed throughout cardiac muscle cells, with predominantly cytoplasmic signal and scattered areas of stronger staining. This RBPMs Antibody demonstrates detection of RNA-Binding Protein With Multiple Splicing in human cardiac muscle tissue.



RBPMs Antibody Mouse Heart IHC. IHC staining of FFPE mouse heart tissue was performed with RBPMs Antibody at 2 ug/ml following heat-induced epitope retrieval in pH 8 EDTA buffer. HRP-conjugated secondary antibody and DAB were used for detection. RBPMs staining is observed in cardiac muscle cells, with predominantly cytoplasmic signal. This RBPMs Antibody demonstrates detection of RNA-Binding Protein With Multiple Splicing in mouse heart tissue.



RBPMs Antibody Rat Heart IHC. IHC staining of FFPE rat heart tissue was performed with RBPMs Antibody at 2 ug/ml following heat-induced epitope retrieval in pH 8 EDTA buffer. HRP-conjugated secondary antibody and DAB were used for detection. RBPMs staining is observed in cardiac muscle cells, with predominantly cytoplasmic signal and stronger staining in scattered cells. This RBPMs Antibody demonstrates detection of RNA-Binding Protein With Multiple Splicing in rat heart tissue.

Description

RBPMs Antibody / RNA-Binding Protein With Multiple Splicing Antibody detects RBPMs, an RNA-binding protein involved in post-transcriptional regulation of gene expression. RBPMs contains a conserved RNA recognition motif that enables interactions with RNA and contributes to the regulation of RNA processing and alternative splicing. The protein is expressed in multiple tissues and has been studied particularly in relation to muscle cell differentiation, cellular identity and the regulation of tissue-specific gene expression.

BPMS has an important role in smooth muscle biology, where it functions as a regulator of alternative splicing programs associated with the differentiated contractile phenotype. Changes in BPMS expression can influence the processing of transcripts involved in cytoskeletal organization, contraction and other characteristics of differentiated muscle cells. A BPMS Antibody can therefore support studies of RNA regulation, muscle differentiation and mechanisms that control transitions between cellular phenotypes.

BPMS expression is also found in cardiac tissue, making the protein relevant to studies of RNA processing and gene regulation in muscle-rich tissues. More broadly, RNA-binding proteins such as BPMS can influence transcript processing, stability and utilization, allowing cells to coordinate protein expression after transcription. These mechanisms are important during development, differentiation and responses to physiological or pathological changes.

In neuroscience, BPMS is also widely used as a marker for retinal ganglion cells (RGCs). BPMS immunoreactivity can be used to identify and quantify RGCs in retinal tissue and has been applied in studies of retinal development, optic nerve injury and neurodegeneration. BPMS is not specific to the retina or retinal ganglion cells, however, and its expression in muscle and other tissues reflects broader biological functions in RNA regulation.

NSJ Bioreagents offers BPMS Antibody reagents for investigating BPMS expression and localization. An RNA-Binding Protein With Multiple Splicing Antibody can support research into RNA processing, alternative splicing, muscle biology, cellular differentiation and retinal ganglion cell biology.

Researchers studying BPMS in retinal ganglion cells and neuronal biology can explore our [Neuroscience Antibodies](#) page for additional targets involved in neural development, function and neurodegeneration.

Application Notes

Optimal dilution of the BPMS Antibody / RNA-Binding Protein With Multiple Splicing Antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids K6-C196) was used as the immunogen for the BPMS antibody.

Storage

After reconstitution, the BPMS antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

Alternate Names

BPMS antibody, RNA-Binding Protein With Multiple Splicing antibody, HERMES antibody, BPMS1 antibody, Retinal Ganglion Cell Marker antibody