


## Recombinant AMACR Antibody / p504S [clone AMACR/2748R] (V7480)

Catalog No.	Formulation	Size
V7480-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7480-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7480SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7480IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG, kappa
<b>Clone Name</b>	AMACR/2748R
<b>Purity</b>	Protein A affinity chromatography
<b>UniProt</b>	Q9UHK6
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 1-2ug/ml
<b>Limitations</b>	This recombinant AMACR antibody is available for research use only.




Recombinant AMACR Antibody RCC IHC. Immunohistochemistry testing of FFPE human renal cell carcinoma with recombinant AMACR antibody. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.




Recombinant AMACR Antibody Prostate Cancer IHC. Immunohistochemistry testing of FFPE human prostate carcinoma with recombinant AMACR antibody. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



SDS-PAGE analysis of purified, BSA-free AMACR antibody (clone AMACR/2748R) as confirmation of integrity and purity.



Western blot testing of human kidney lysate with recombinant AMACR antibody. Predicted molecular weight ~43 kDa.



Recombinant AMACR Antibody Kidney Tissue WB. Western blot analysis of human, mouse, rat, hamster, and guinea pig kidney tissue lysates using AMACR Antibody clone AMACR/2748R demonstrates a distinct band at approximately 40-42 kDa selectively in human kidney lysate, consistent with the predicted molecular weight of Alpha-methylacyl-CoA racemase / AMACR. The observed species-selective reactivity pattern suggests preferential recognition of the human AMACR epitope by clone AMACR/2748R.

## Description

Recombinant AMACR Antibody recognizes a protein of 43kDa, which is identified as AMACR, also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and -oxidation of branched-chain fatty acids. AMACR is essential in lipid metabolism. It is expressed in cells of premalignant high-grade prostatic intraepithelial neoplasia (HGPIN) and prostate adenocarcinoma. The majority of the carcinoma cells show a distinct granular cytoplasmic staining reaction. AMACR is present at low or undetectable levels in glandular epithelial cells of normal prostate and benign prostatic hyperplasia. A spotty granular cytoplasmic staining is seen in a few cells of the benign glands. AMACR is expressed in normal liver (hepatocytes), kidney (tubular epithelial cells) and gall bladder (epithelial cells). Expression has also been found in lung (bronchial epithelial cells) and colon (colonic surface epithelium). AMACR expression can also be found in hepatocellular carcinoma and kidney carcinoma. Past studies have also shown that AMACR is expressed in various colon carcinomas (well, moderately and poorly differentiated) and over expressed in prostate carcinoma.

For additional AMACR and p504S research antibodies validated by protein microarray specificity analysis, western blotting, and immunohistochemistry, explore the broader [AMACR Antibody page](#) featuring clone AMACR/1864.

## Application Notes

Optimal dilution of the recombinant AMACR antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## **Immunogen**

A portion of amino acids 297-394 was used as the immunogen for the recombinant AMACR antibody.

## **Storage**

Store the recombinant AMACR antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).