


Recombinant Anaplastic Lymphoma Kinase Antibody / ALK [clone ALK1/9660R] (V5626)

Catalog No.	Formulation	Size
V5626-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5626-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5626SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days	
Species Reactivity	Human	
Format	Purified	
Clonality	Recombinant Rabbit Monoclonal	
Isotype	Rabbit IgG, kappa	
Clone Name	ALK1/9660R	
Purity	Protein A/G affinity	
UniProt	Q9UM73	
Localization	Cell membrane	
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml	
Limitations	This recombinant Anaplastic Lymphoma Kinase antibody is available for research use only.	



IHC staining of FFPE human Ewing Sarcoma tissue with recombinant Anaplastic Lymphoma Kinase antibody (clone ALK1/9660R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Description

Anaplastic lymphoma kinase (ALK) is a receptor tyrosine kinase of the insulin receptor superfamily. ALK is typically expressed at low levels in regions of the developing central and peripheral nervous system. ALK may be activated in cancer through multiple mechanisms. The most common mechanism is through formation of a fusion protein from chromosomal translocations, as in the case of anaplastic large cell lymphoma (ALCL) and inflammatory myofibroblastic tumors. ALK may also be amplified through mutation, as in neuroblastomas. Various solid tumors, such as non-small cell lung carcinoma (NSCLC) and brain cancers were also found to aberrantly express ALK. ALK staining is present within both the nucleus and cytoplasm, and are positive in about 60% of ALCL. ALK protein expression by tumor cells is an independent prognostic factor that predicts a favorable outcome.

Application Notes

Optimal dilution of the recombinant Anaplastic Lymphoma Kinase antibody should be determined by the researcher.

Immunogen

Recombinant human ALK protein fragment corresponding to the cytoplasmic domain of the protein was used as the immunogen for the recombinant Anaplastic Lymphoma Kinase antibody.

Storage

Aliquot the recombinant Anaplastic Lymphoma Kinase antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.