


Recombinant Bromodeoxyuridine Antibody / BrdU [clone rBRD469] (V8337)

| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V8337-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8337-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V8337SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | All species |
| Format | Purified |
| Host | Mouse |
| Clonality | Recombinant Mouse Monoclonal |
| Isotype | Mouse IgG1, kappa |
| Clone Name | rBRD469 |
| Purity | Protein G affinity chromatography |
| Localization | Nuclear |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This recombinant Bromodeoxyuridine antibody is available for research use only. |



SDS-PAGE analysis of purified, BSA-free recombinant Bromodeoxyuridine antibody (clone rBRD469) as confirmation of integrity and purity.

Description

It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei

during DNA synthesis prior to mitosis. Antibody to BrdU is helpful in detecting S-phase cells, providing useful information on the aggressiveness of tumors.

Application Notes

Optimal dilution of the recombinant Bromodeoxyuridine antibody should be determined by the researcher.

Immunogen

Bromodeoxyuridine conjugated to KLH was used as the immunogen.

Storage

Store the recombinant Bromodeoxyuridine antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).