


Recombinant Cadherin 1 Antibody / CDH1 / E-Cadherin [clone rCDH1/6769] (V5471)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V5471-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V5471-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V5471SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |


Recombinant **MOUSE MONOCLONAL**

Bulk quote request


| | |
|--------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Recombinant Mouse Monoclonal |
| Isotype | Mouse IgG1, kappa |
| Clone Name | rCDH1/6769 |
| Purity | Protein A/G affinity |
| UniProt | P12830 |
| Localization | Cell membrane |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This recombinant Cadherin 1 antibody is available for research use only. |



IHC staining of FFPE human stomach tissue with recombinant Cadherin 1 antibody (clone rCDH1/6769). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human colon tissue with recombinant Cadherin 1 antibody (clone rCDH1/6769). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Cadherin 1 antibody (rCDH1/6769) as confirmation of integrity and purity.

Description

E-cadherin is a transmembrane, calcium dependent cell adhesion protein that mediates cell to cell adhesion and maintains structural and functional integrity of epithelial tissues. It also has pivotal barrier functions and maintains the polarity of epithelial cells. Reduced or aberrant E-cadherin expression breaks cell to cell contacts, and thus, cells acquire the ability to migrate. In normal tissues, immunostaining of E-cadherin is localized to the membrane of epithelial cells, consistent with its role in cell adhesion. And in tumor tissues, E-cadherin stains positively in glandular epithelium as well as adenocarcinomas of the lung, gastrointestinal tract, and ovary. It has also been shown to be positive in some thyroid carcinomas. A combination of E-cadherin and p120 catenin may help distinguish ductal carcinoma of the breast from lobular carcinoma. And also, several studies have reported that reduced E-cadherin expression is correlated with poor prognosis in several types of carcinomas.

Application Notes

Optimal dilution of the recombinant Cadherin 1 antibody should be determined by the researcher.

Immunogen

A recombinant fragment (within amino acids 600-700) of human Cadherin 1 was used as the immunogen for the recombinant Cadherin 1 antibody.

Storage

Aliquot the recombinant Cadherin 1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.