


## Recombinant Carbonic Anhydrase 9 Antibody [clone CA9/2993R] (V7364)

Catalog No.	Formulation	Size
V7364-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7364-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7364SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7364IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG, kappa
<b>Clone Name</b>	CA9/2993R
<b>Purity</b>	Protein A affinity chromatography
<b>UniProt</b>	Q16790
<b>Localization</b>	Cell surface and cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
<b>Limitations</b>	This recombinant Carbonic Anhydrase 9 antibody is available for research use only.



IHC staining of FFPE human renal cell carcinoma with recombinant Carbonic Anhydrase 9 antibody (clone CA9/2993R). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Carbonic Anhydrase 9 antibody (clone CA9/2993R) as confirmation of integrity and purity.

## Description

Recognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowman's capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This MAb may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.

## Application Notes

The optimal dilution of the recombinant Carbonic Anhydrase 9 antibody for each application should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

Recombinant human protein was used as the immunogen for this recombinant Carbonic Anhydrase 9 antibody.

## Storage

Store the recombinant Carbonic Anhydrase 9 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).