


## Recombinant CCNE1 Antibody / Cyclin E1 [clone CCNE1/4935R] (V9178)

Catalog No.	Formulation	Size
V9178-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9178-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9178SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Name</b>	CCNE1/4935R
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	IDP24864
<b>Localization</b>	Nuclear
<b>Applications</b>	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This recombinant CCNE1 antibody is available for research use only.




Western blot testing of human HCT-116 cell lysate using recombinant CCNE1 antibody (clone CCNE1/4935R). Predicted molecular weight ~47 kDa.




IHC staining of FFPE human endometrium with recombinant CCNE1 antibody (clone CCNE1/4935R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human ovarian carcinoma tissue with recombinant CCNE1 antibody (clone CCNE1/4935R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant CCNE1 antibody (clone CCNE1/4935R) as confirmation of integrity and purity.



Western blot testing of human HCT-116 cell lysate using recombinant CCNE1 antibody (clone CCNE1/4935R). Predicted molecular weight ~47 kDa.

## Description

Cyclin E belongs to the highly conserved cyclin family, whose members exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. Cyclins function as regulators of CDK kinases. Cyclin E forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. Cyclin E accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Cyclin E overexpression has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis.

## Application Notes

Optimal dilution of the recombinant CCNE1 antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 10-176 was used as the immunogen for the recombinant CCNE1 antibody.

## Storage

Aliquot the recombinant CCNE1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

