


Recombinant CD146 Antibody [clone rMUC18/1130] (V3638)

Catalog No.	Formulation	Size
V3638-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3638-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3638SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3638IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rMUC18/1130
Purity	Protein G affinity chromatography
UniProt	P43121
Localization	Cell surface, cytoplasm
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This recombinant CD146 antibody is available for research use only.



IHC testing of FFPE human tonsil with CD146 antibody (clone rMUC18/1130). Required boiling tissue sections in 10mM Tris with 1mM EDTA, pH8, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant CD146 antibody (clone rMUC18/1130) as confirmation of integrity and purity.

Description

CD146 expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not normal melanocytes. It is found on 80% of advanced primary human melanomas and correlates well with development of metastatic disease.

Application Notes

Optimal dilution of the recombinant CD146 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the recombinant CD146 antibody.

Storage

Store the recombinant CD146 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).