


Recombinant CD40 Antibody [clone C40/9225R] (V5525)

Catalog No.	Formulation	Size
V5525-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5525-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5525SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG1, kappa
Clone Name	C40/9225R
Purity	Protein A/G affinity
UniProt	P25942
Localization	Cell Surface, Secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant CD40 antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with recombinant CD40 antibody (clone C40/9225R). Inset: PBS used in place of primary Ab (secondary Ab negative control).
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant CD40 antibody (clone C40/9225R) as confirmation of integrity and purity.

Description

CD40 is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. CD40 is expressed on B-lymphocytes, follicular dendritic cells, bone marrow-derived dendritic cells, thymic epithelium, and interdigitating cells in the T-cell zones of secondary lymphoid organs.

Application Notes

Optimal dilution of the recombinant CD40 antibody should be determined by the researcher.

Immunogen

Recombinant full-length human CD40 protein was used as the immunogen for the recombinant CD40 antibody.

Storage

Aliquot the recombinant CD40 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.