
Recombinant CD45 Antibody PTPRC/1975R + PTPRC/1783R [clone PTPRC/1975R + PTPRC/1783R] (V7330)

Catalog No.	Formulation	Size
V7330-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7330-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7330SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7330IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	PTPRC/1975R + PTPRC/1783R
Purity	Protein A affinity chromatography
UniProt	P08575
Localization	Cell surface and cytoplasmic
Applications	Western blot: 1-2ug/ml Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Limitations	This Recombinant CD45 Antibody PTPRC/1975R + PTPRC/1783R is available for research use only.



Recombinant CD45 Antibody PTPRC/1975R + PTPRC/1783R Immunohistochemistry. IHC testing of recombinant CD45 antibody cocktail and FFPE human tonsil (clones PTPRC/1975R + PTPRC/1783R). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.



Recombinant CD45 Antibody PTPRC/1975R + PTPRC/1783R WB. Western blot testing of human spleen lysate with recombinant CD45 antibody.

Description

Recombinant CD45 Antibody PTPRC/1975R + PTPRC/1783R detects CD45R, also designated CD45 and PTPRC, identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.

CD45 is encoded by PTPRC and functions as a key regulator of leukocyte signaling and hematopoietic cell biology; additional reagents targeting this protein are available on the CD45 Antibody page.

Application Notes

Due to variation in protocol and secondary antibody used, the Recombinant CD45 Antibody PTPRC/1975R + PTPRC/1783R may need to be titered for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as immunogen for this recombinant CD45 antibody cocktail.

Storage

The recombinant CD45 antibody cocktail (with azide) can be stored at 2-8oC. The azide-free format should be aliquoted and stored at -20oC or colder.

References (2)

Copyright © NSJ Bioreagents. All rights reserved.