


Recombinant CD54 Antibody / ICAM-1 [clone ICAM1/8677R] (V4610)

Catalog No.	Formulation	Size
V4610-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4610-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4610SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	ICAM1/8677R
Purity	Protein A/G affinity
UniProt	P05362
Localization	Cell Surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This recombinant CD54 antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with recombinant CD54 antibody (clone ICAM1/8677R). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant CD54 antibody (clone ICAM1/8677R) as confirmation of integrity and purity.

Description

ICAM-1 is a single chain glycoprotein of Ig supergene family, present on unstimulated endothelial cells (EC) and on a variety of other cell types including activated fibroblasts, EC, macrophages, and lymphocytes.

Application Notes

Optimal dilution of the recombinant CD54 antibody should be determined by the researcher.

Immunogen

A recombinant fragment corresponding to the intracellular portion of the human protein was used as the immunogen for the recombinant CD54 antibody.

Storage

Aliquot the recombinant CD54 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.