


# Recombinant CD59 Antibody [clone MACIF/2867R] (V7387)

Catalog No.	Formulation	Size
V7387-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7387-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7387SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7387IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


# Recombinant RABBIT MONOCLONAL

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MACIF/2867R
Purity	Protein A affinity
UniProt	P13987
Localization	Plasma membrane, cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This recombinant CD59 antibody is available for research use only.



IHC staining of FFPE human colon carinoma with recombinant CD59 antibody (clone MACIF/2867R) . HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant CD59 antibody (clone MACIF/2867R) as confirmation of integrity and purity.

#### **Description**

Reacts with human CD59, a 20kDa glycosyl phosphatidyl-inositol (GPI)-anchored cell surface protein. CD59 regulates complement-mediated cell lysis, and it is involved in lymphocyte signal transduction. This protein is a potent inhibitor of the complement membrane attack complex, whereby it binds complement C8 and/or C9 during the assembly of this complex, thereby inhibiting the incorporation of multiple copies of C9 into the complex, which is necessary for osmolytic pore formation. CD59 is widely distributed on cells in all tissues. It inhibits formation of MAC, thus protecting cells from complement-mediated lysis. The expression of CD59 on erythrocytes is important for their survival. Genetic defects in GPI-anchor attachment, that cause a reduction or loss of CD59 and CD55 on erythrocytes produce the symptoms of the disease paroxysmal hemoglobinuria (PNH). It is useful for study on GPI-anchored proteins, PNH and CD59 functions.

## **Application Notes**

Optimal dilution of the recombinant CD59 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

Recombinant full-length human protein was used as the immunogen for the recombinant CD59 antibody.

#### **Storage**

Store the recombinant CD59 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).