


# Recombinant CD61 Antibody / Integrin beta 3 / ITGB3 [clone ITGB3/3126R] (V8142)

Catalog No.	Formulation	Size
V8142-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8142-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8142SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


## Recombinant RABBIT MONOCLONAL

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	ITGB3/3126R
Purity	Protein A affinity chromatography
UniProt	P05106
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant CD61 antibody is available for research use only.



IHC testing of FFPE human spleen with recombinant CD61 antibody (ITGB3/3126R). Required HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH9, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant CD61 antibody (clone ITGB3/3126R) as confirmation of integrity and purity.

#### **Description**

Reacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.

#### **Application Notes**

Optimal dilution of the recombinant CD61 antibody should be determined by the researcher.

#### **Immunogen**

A recombinant human partial protein (amino acids 385-490) was used as the immunogen for this recombinant CD61 antibody.

#### **Storage**

Store the recombinant CD61 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).