


Recombinant CD63 Antibody / LAMP-3 [clone rMX-49.129.5] (V7388)

Catalog No.	Formulation	Size
V7388-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7388-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7388SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7388IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rMX-49.129.5
Purity	Protein G affinity chromatography
Gene ID	967
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Flow Cytometry : 1-2ug/million cells
Limitations	This recombinant CD63 antibody is available for research use only.



Flow cytometry testing of PFA-fixed human U-87 MG cells with recombinant CD63 antibody (clone rMX-49.129.5); Red=isotype control, Blue= CD63 antibody.



SDS-PAGE analysis of purified, BSA-free recombinant CD63 antibody (clone rMX-49.129.5) as confirmation of integrity and purity.



IHC staining of FFPE human melanoma with recombinant CD63 antibody (clone rMX-49.129.5). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using recombinant CD63 antibody (clone rMX-49.129.5). These results demonstrate the foremost specificity of the rMX-49.129.5 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

Application Notes

Immunogen

Full length human CD63 was used as the immunogen for this recombinant CD63 antibody.

Storage

Store the recombinant CD63 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

References (3)

