


Recombinant CD7 Antibody [clone rCD7/6387] (V9149)

Catalog No.	Formulation	Size
V9149-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9149-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9149SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2a, kappa
Clone Name	rCD7/6387
Purity	Protein A/G affinity
UniProt	P09564
Localization	Cell Surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant CD7 antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with recombinant CD7 antibody (clone rCD7/6387). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Negative control: IHC staining of FFPE human brain tissue using recombinant CD7 antibody (clone rCD7/6387) at 2ug/ml in PBS for 30min RT. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant CD7 antibody (clone rCD7/6387) as confirmation of integrity and purity.

Description

Recognizes a protein of 40kDa, identified as CD7 (also known as gp40, Leu9). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

Application Notes

Optimal dilution of the recombinant CD7 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 72-175 was used as the immunogen for the recombinant CD7 antibody.

Storage

Aliquot the recombinant CD7 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.