


## Recombinant CD71 Antibody / Transferrin Receptor / TFRC (Extracellular domain) [clone TFRC/2898R] (V8115)

Catalog No.	Formulation	Size
V8115-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8115-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8115SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

**Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	TFRC/2898R
Purity	Protein A affinity chromatography
UniProt	P02786
Localization	Cell surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant CD71 antibody is available for research use only.



IHC staining of FFPE human bone marrow with recombinant CD71 antibody (clone TFRC/2898R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant CD71 antibody antibody (clone TFRC/2898R) as confirmation of integrity and purity.

## Description

It recognizes a ~90-95kDa protein which is identified as cell surface transferrin receptor (CD71), a disulfide-bonded homodimeric glycoprotein of 180-190kDa. This MAb is highly specific to CD71 and shows no cross-reaction with other related proteins. Ligand for transferrin receptor is the serum iron transport protein, transferrin. This receptor is broadly distributed in carcinomas, sarcomas, leukemias, and lymphomas. CD71/Transferrin receptor has been reported to be associated with cell proliferation in both normal and neoplastic tissues and useful in predicting clinical behavior or response to therapy in a number of malignancies including breast cancer.

## Application Notes

Optimal dilution of the recombinant CD71 antibody should be determined by the researcher.

## Immunogen

A recombinant human partial protein (extracellular region, amino acids 94-212) was used as the immunogen for this recombinant CD71 antibody.

## Storage

Store the recombinant CD71 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).