


## Recombinant CD79a Antibody / Rabbit Monoclonal [clone CDLA79a-3R] (V3696)

Catalog No.	Formulation	Size
V3696-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3696-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3696SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3696IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Recombinant RABBIT MONOCLONAL

**Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	CDLA79a-3R
Purity	Protein A affinity chromatography
UniProt	P11912
Localization	Cell surface, cytoplasmic
Applications	Western Blot : 1-2ug/ml Immunofluorescence : 1-2ug/ml Flow Cytometry : 1-2ug/10 <sup>6</sup> cells Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT (1)
Limitations	This recombinant CD79a antibody is available for research use only.




Flow cytometry testing of human Raji cells with recombinant CD79a antibody (clone CDLA79a-3R); Red=isotype control, Blue= CD79a antibody.



Western blot testing of human Raji cell lysate with recombinant CD79a antibody (clone CDLA79a-3R). Expected molecular weight: 25-47 kDa depending on glycosylation level.



Immunofluorescent staining of PFA-fixed human Raji cells with recombinant CD79a antibody (green, clone CDLA79a-3R) and Reddot nuclear stain (red).



IHC testing of FFPE human tonsil with recombinant CD79a antibody (clone CDLA79a-3R). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA followed by cooling at RT for 20 min.

## Description

The CD79a protein together with the related CD79b protein, forms a dimer associated with membrane-bound immunoglobulin in B-cells, thus forming the B-cell antigen receptor (BCR). CD79a plays multiple and diverse roles in B cell development and function. The CD79a/b heterodimer associates non-covalently with the immunoglobulin heavy chain through its transmembrane region, thus forming the BCR along with the immunoglobulin light chain and the pre-BCR when associated with the surrogate light chain in developing B cells. Association of the CD79a/b heterodimer with the immunoglobulin heavy chain is required for surface expression of the BCR and BCR induced calcium flux and protein tyrosine phosphorylation. [Wiki]

## Application Notes

The stated application concentrations are suggested starting points. Titration of the CD79a antibody may be required due to differences in protocols and secondary/substrate sensitivity.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

Amino acids 202-216 (GTYQDVGSNIADVQ) were used as the immunogen for the recombinant CD79a antibody.

## Storage

Store the recombinant CD79a antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).