
Recombinant CDX2 Antibody [clone CDX2/2951R] (V3964)

Catalog No.	Formulation	Size
V3964-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3964-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3964SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3964IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant RABBIT MONOCLONAL

Bulk quote request


Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	CDX2/2951R
Purity	Protein A affinity chromatography
UniProt	Q99626
Localization	Nuclear
Applications	ELISA : 2-4ug/ml (order BSA/azide-free format) Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 2-4ug/ml
Limitations	This recombinant CDX2 antibody is available for research use only.




Recombinant CDX2 Antibody Colon Carcinoma IHC. Immunohistochemistry testing of FFPE human colon carcinoma with recombinant CDX2 antibody (clone CDX2/2951R). HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling prior to testing.



Recombinant CDX2 Antibody Human Colon Carcinoma Tissue IHC. Immunohistochemistry testing of FFPE human colon carcinoma with recombinant CDX2 antibody (clone CDX2/2951R). HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling prior to testing.



SDS-PAGE analysis of purified, BSA-free recombinant CDX2 antibody (clone CDX2/2951R) as confirmation of integrity and purity.



CDX2 Antibody Human/Cell Line WB. Western blot analysis of CDX2 / Caudal type homeobox 2 expression using CDX2 antibody, clone CDX2/2951R. Lane 1: HEK293T cell lysate, Lane 2: human colon tissue lysate, Lane 3: human small intestine tissue lysate. A band is detected at approximately 40 kDa, consistent with the predicted molecular weight of CDX2, with stronger signal observed in intestinal tissues compared to HEK293T cells, aligning with the known epithelial and gastrointestinal-restricted expression pattern of this transcription factor.

Description

Recombinant CDX2 Antibody detects Caudal-type homeobox protein 2. The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CDX2 has been useful to establish GI origin of metastatic adenocarcinomas and carcinoids and is especially useful to distinguish metastatic colorectal adenocarcinoma from lung adenocarcinoma. However, mucinous carcinomas of the ovary also express CDX2 protein. It limits the usefulness of this marker in the distinction of metastatic colorectal adenocarcinoma from mucinous carcinoma of the ovary.

This antibody is part of a broader antibody panel offered by NSJ Bioreagents.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the recombinant CDX2 antibody to be titered up or down for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 150-249 from the human protein were used as the immunogen for this recombinant CDX2 antibody.

Storage

Store the recombinant CDX2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

Caudal type homeo box 2 antibody, Caudal type homeo box transcription factor 2 antibody

Copyright © NSJ Bioreagents. All rights reserved.