


## Recombinant CFTR Antibody / Cystic Fibrosis Transmembrane Regulator [clone rCFTR/7175] (V9529)

| Catalog No.    | Formulation   | Size   |
|----------------|---|--------|
| V9529-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V9529-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug  |
| V9529SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                          | 100 ug |

### Recombinant MOUSE MONOCLONAL

[Bulk quote request](#)

|                           |  |
|---------------------------|--|
| <b>Availability</b>       | 1-3 business days  |
| <b>Species Reactivity</b> | Human  |
| <b>Format</b>             | Purified   |
| <b>Host</b>               | Mouse  |
| <b>Clonality</b>          | Recombinant Mouse Monoclonal                                       |
| <b>Isotype</b>            | Mouse IgG1, kappa  |
| <b>Clone Name</b>         | rCFTR/7175   |
| <b>Purity</b>             | Protein A/G affinity   |
| <b>UniProt</b>            | P13569   |
| <b>Localization</b>       | Cell surface, Cytoplasm  |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 1-2ug/ml                             |
| <b>Limitations</b>        | This recombinant CFTR antibody is available for research use only. |



IHC staining of FFPE human salivary gland tissue with recombinant CFTR antibody (clone rCFTR/7175) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is

composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation.

For a [widely cited CFTR antibody](#) used across multiple study types, see our M3A7 clone.

## Application Notes

Optimal dilution of the recombinant CFTR antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 550-850 was used as the immunogen for the recombinant CFTR antibody.

## Storage

Aliquot the recombinant CFTR antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.