


Recombinant CTNND1 Antibody / p120 Catenin / Catenin delta 1 [clone rCTNND1/6903] (V9528)

Catalog No.	Formulation	Size
V9528-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9528-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9528SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant MOUSE MONOCLONAL

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rCTNND1/6903
Purity	Protein A/G affinity
UniProt	O60716
Localization	Cell Surface, Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant CTNND1 antibody is available for research use only.



IHC staining of FFPE human stomach tissue glandular cells with recombinant CTNND1 antibody (clone rCTNND1/6903) at 2ug/ml in PBS for 30min RT. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant CTNND1 antibody (clone rCTNND1/6903) as confirmation of integrity and purity.

Description

Alpha-catenin and beta-catenin bind to the intracellular domain of E-cadherin while p120 catenin binds E-cadherin at a juxta-membrane site. The complex stabilizes tight junctions. In the cell, p120 catenin localized to the E-cadherin/catenins cell adhesion complex, directly associates with cytoplasmic C-terminus of E-cadherin and may similarly interact with other cadherins. p120 is a proliferation-associated nucleolar protein found in most human malignant tumors, but not in resting normal cells. In colorectal cancer the altered localization of p120 catenin corresponds with loss of cytoplasmic localization of E-cadherin. Studies have shown accurate categorization of ductal vs. lobular neoplasia in the breast was achieved with p120 staining. p120 expression further clarifies the separation of low-grade ductal carcinoma in situ from lobular neoplasia. Studies also have shown that altered expression of p120 catenin antibody predicts poor outcome in invasive breast cancer.

Application Notes

Optimal dilution of the recombinant CTNND1 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 868-968 was used as the immunogen for the recombinant CTNND1 antibody.

Storage

Aliquot the recombinant CTNND1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.