


Cytokeratin 19 Antibody / CK19 / Rabbit Monoclonal [clone CTKN19-2R] (V3743)

Catalog No.	Formulation	Size
V3743-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3743-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3743SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3743IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	CTKN19-2R
Purity	Protein A affinity chromatography
UniProt	P08727
Localization	Cytoplasmic, cell membrane, secreted
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Prediluted IHC Only Format : incubate for 30 min at RT (1)
Limitations	This recombinant Cytokeratin 19 antibody is available for research use only.



Western blot testing of human HepG2 antibody with Cytokeratin 19 antibody (clone CTKN19-2R). Predicted molecular weight ~43 kDa.



Immunohistochemistry analysis of recombinant Cytokeratin 19 / CK19 antibody (clone CTKN19-2R) in human colon tissue. Formalin-fixed, paraffin-embedded human colon tissue was stained using recombinant Cytokeratin 19 antibody (clone CTKN19-2R). Heat-induced epitope retrieval was performed by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes, followed by cooling at room temperature prior to antibody incubation. Brown chromogenic signal is observed in glandular epithelial cells lining colonic crypts, with strong cytoplasmic and membranous staining outlining epithelial structures, while surrounding stromal tissue shows little to no staining. This staining pattern reflects epithelial-associated expression of Cytokeratin 19 in colonic epithelium.

Description

Cytokeratin 19 Antibody recognizes Cytokeratin 19, also known as Keratin 19 (KRT19), a type I intermediate filament protein that is widely expressed in simple epithelial cells and in progenitor-associated epithelial compartments of stratified tissues. Cytokeratin 19 is the smallest member of the type I keratin family and lacks the C-terminal tail domain present in most keratins, a structural feature that influences its filament organization and cellular distribution. Cytokeratin 19 Antibody is commonly used in research and pathology contexts and is frequently referred to in the literature as CK19 antibody or Keratin 19 antibody.

Cytokeratin 19 expression is characteristic of epithelial cells lining glandular and ductal structures, including bile ducts, pancreatic ducts, renal tubules, bronchial epithelium, gastrointestinal mucosa, mammary epithelium, and prostatic glands. In stratified epithelia, CK19 expression is often enriched in basal or progenitor-associated epithelial cells rather than terminally differentiated suprabasal layers. This expression pattern distinguishes Cytokeratin 19 from differentiation-associated keratins such as Cytokeratin 10 or Cytokeratin 13 and makes Cytokeratin 19 Antibody useful for identifying epithelial cells with ductal, glandular, or progenitor-like phenotypes.

Alterations in Cytokeratin 19 expression have been reported in a variety of pathological contexts. Changes in CK19 expression patterns are frequently observed in epithelial-derived malignancies, including carcinomas of the breast, lung, pancreas, thyroid, liver, and gastrointestinal tract. As a result, Cytokeratin 19 antibody staining patterns are commonly evaluated in research studies focused on epithelial differentiation, ductal lineage identification, and carcinoma biology, particularly in tumors exhibiting glandular or ductal differentiation.

At the cellular level, Cytokeratin 19 contributes to the organization of the intermediate filament cytoskeleton and supports epithelial cell integrity and structural adaptability. Its broad expression in simple and ductal epithelia makes Cytokeratin 19 Antibody a valuable tool for studies of epithelial lineage tracing, tissue architecture, and epithelial biology. The Cytokeratin 19 Antibody (clone CTKN19-2R) is designed to detect Cytokeratin 19 expression in research applications where assessment of epithelial and ductal cell populations is required.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant Cytokeratin 19 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if

required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Human mammary epithelial organoids were used as the immunogen for the recombinant Cytokeratin 19 antibody.

Storage

Store the recombinant Cytokeratin 19 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).