


Recombinant Cytokeratin 5 Antibody [clone KRT5/4245R] (V8736)

Catalog No.	Formulation	Size
V8736-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8736-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8736SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	KRT5/4245R
Purity	Protein A affinity chromatography
UniProt	P13647
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This recombinant Cytokeratin 5 antibody is available for research use only.



IHC staining of FFPE human basal cell carcinoma with recombinant Cytokeratin 5 antibody (clone KRT5/4245R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human tonsil with recombinant Cytokeratin 5 antibody (clone KRT5/4245R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

This MAb recognizes a protein of 58kDa, which is identified as Cytokeratin 5 (KRT5). This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Antibodies to KRT5 identify basal cells of squamous and glandular epithelia, myoepithelia, and mesothelium. Anti-cytokeratin 5 has been reported useful in the differential diagnosis of metastatic carcinoma in the pleura versus epithelioid mesothelioma. Almost all squamous cell carcinomas, half of transitional carcinomas, and many undifferentiated large cell carcinomas express. Anti-KRT5, along with anti-p63, affords a high sensitivity and specificity for squamous differentiation. Myoepithelial cells of the breast, glandular epithelia, and basal cells of the prostate are labeled with anti-KRT5.

Application Notes

Optimal dilution of the recombinant Cytokeratin 5 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 490-590 corresponding to C-terminal region of the human protein was used as the immunogen for the recombinant Cytokeratin 5 antibody.

Storage

Store the recombinant Cytokeratin 5 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).