


Recombinant EGF Receptor / EGFR Antibody (Extracellular domain) [clone rEGFR/6389] (V9273)

Catalog No.	Formulation	Size
V9273-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9273-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9273SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant MOUSE MONOCLONAL

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2a, kappa
Clone Name	rEGFR/6389
Purity	Protein A/G affinity
UniProt	P00533
Localization	Cell Surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant EGF Receptor antibody is available for research use only.



IHC staining of FFPE human placental tissue with recombinant EGF Receptor antibody (clone rEGFR/6389). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Negative control: IHC staining of FFPE human testis tissue with recombinant EGF Receptor antibody (clone rEGFR/6389) at 2ug/ml in PBS for 30min RT. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant EGF Receptor antibody (clone rEGFR/6389) as confirmation of integrity and purity.

Description

This MAbs recognizes a protein of 170kDa, identified as EGFR. EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor- α (TGF- α), Heparin-binding EGF (HB-EGF), amphiregulin, betacellulin and epiregulin. EGFR is overexpressed in tumors of breast, brain, bladder, lung, gastric, head neck, esophagus, cervix, vulva, ovary, and endometrium. It is predominantly present in squamous cell carcinomas.

Application Notes

Optimal dilution of the recombinant EGF Receptor antibody should be determined by the researcher.

Immunogen

Amino acids 25-645 from the extracellular domain of the human protein were used as the immunogen for the recombinant EGF Receptor antibody.

Storage

Aliquot the recombinant EGF Receptor antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.