


Recombinant FLG Antibody / Filaggrin [clone FLG/1957R] (V3612)

Catalog No.	Formulation	Size
V3612-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3612-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3612SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3612IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	FLG/1957R
Purity	Protein G affinity chromatography
UniProt	P20930
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This recombinant FLG antibody is available for research use only.



IHC testing of FFPE human skin with recombinant FLG antibody (clone FLG/1957R). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant FLG antibody (clone FLG/1957R) as confirmation of integrity and purity.

Description

Filaggrin (FLG) is an intermediate filament-associated protein that aggregates keratin intermediate filaments in mammalian epidermis. It is initially synthesized as a polypeptide precursor, profilaggrin (consisting of multiple Filaggrin units of 324 aa each), which is localized in keratohyalin granules, and is subsequently proteolytically processed into individual functional Filaggrin molecules. Active Filaggrin is present at a level of the epidermis where keratinocytes are in transition between the live nucleated granular layer and the anucleate cornified layer, suggesting that filaggrin aids in the terminal differentiation process by facilitating apoptotic machinery.

Application Notes

Optimal dilution of the recombinant FLG antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 198-288 of human Filaggrin were used as the immunogen for this recombinant FLG antibody.

Storage

Store the recombinant FLG antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).