


# Recombinant H3K4me3 Antibody [clone RM137] (R20202)

Catalog No.	Formulation	Size
R20202-100UG	1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide	100 ug
R20202-25UG	1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide	25 ug


## Recombinant RABBIT MONOCLONAL

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	All Species
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	RM137
Purity	Protein A purified from animal origin-free supernatant
UniProt	P84243
Gene ID	8350
Applications	Western Blot: 0.2-1ug/ml ELISA: 0.1-0.5ug/ml Dot Blot: 0.1-0.5ug/ml
Limitations	This recombinant H3K4me3 antibody is available for research use only.



Western blot of recombinant histone H3.3 (1) and acid extracts of human HeLa cells (2) using the recombinant H3K4me3 antibody at 0.5 ug/ml showed a band of Histone H3 trimethylated at Lysine 4 (K4me3) in HeLa cells.



A peptide dot blot shows the recombinant H3K4me3 antibody reacts only to Histone H3 trimethyl-Lysine 4 (K4me3). No cross reactivity with nonmodified (H3N1-19), monomethylated (K4me1) or dimethylated Lysine 4 (K4me2).



The recombinant H3K4me3 antibody specifically reacts to Histone H3 trimethylated at Lysine 4 (K4me3). No cross reactivity with other methylated lysines in Histone H3.

## **Description**

This recombinant H3K4me3 antibody reacts to Histone H3 trimethylated at Lysine 4 (K4me3). No cross reactivity with monomethylated (K4me1) or dimethylated (K4me3), or other methylation in Histone H3.

#### **Application Notes**

The stated application concentrations are suggested starting points. Titration of the recombinant H3K4me3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

### **Immunogen**

A trimethyl-peptide corresponding to Trimethyl-Histone H3 (Lys4) was used as the immunogen for this recombinant H3K4me3 antibody.

#### **Storage**

Store the recombinant H3K4me3 antibody at -20oC (with glycerol) or aliquot and store at -20oC (without glycerol).